## **CLAIMS**

1. L-(-)-moprolol L-(+)-tartrate salt (2:1).

5

10

- 2. Pharmaceutical composition for ophthalmic use, characterized in that it comprises L-(-)-moprolol L-(+)-tartrate (2:1) together with at least one pharmaceutically acceptable vehicle.
- Pharmaceutical composition according to Claim 2, characterized in that it is in the form of a gel, an ointment or eyedrops.
- Pharmaceutical composition according to Claim 2 or 3, characterized in that the amount of L-(-)-moprolol is between 0.01% and 20% by weight.
- Pharmaceutical composition according to Claim 2 or 3, characterized in that the amount of L-(-)-moprolol is between 1% and 8% by weight.
- Process for preparing L-(-)-moprolol L-(+)-tartrate (2:1),
  characterized in that it includes the addition of L-(+)-tartaric acid, dissolved in a suitable organic solvent, to L-(-)-moprolol base, also dissolved in a suitable organic solvent, in a 2:1 molar ratio.
  - 7. Process according to Claim 6, characterized in that the salt thus formed is isolated via precipitation and filtration.
- 20 8. Process according to Claim 6 or 7, characterized in that the abovementioned organic solvent is ethyl alcohol.
  - Process according to Claim 8, characterized in that the salt is precipitated from the ethanolic solution via addition of ethyl ether.